

PBA vs. Depression

Onscreen Text:

PBA: a neurologic disorder that can be mistaken for depression

While PBA and depression are commonly comorbid, **PBA is not a mood disorder**. It's a **distinct neurologic condition caused by** other neurologic conditions.

Onscreen Text:

Sequena

Living with PBA

Sequena: My emotions weren't matching what my body was, you know, saying or doing or feeling it made me feel like something was wrong.

Onscreen Text:

Jill

Living with PBA

Jill: Sometimes I would find myself with an outburst of crying just while I was grocery shopping or doing something very mundane and not feeling particularly depressed about anything or upset.

Sequena: I was confused because they were telling me that I had depression. They were prescribing me antidepressants, but yet I was still crying all the time. It had to be something else.

Onscreen Text:

For your patients with **PBA**

NUEDEXTA CAN MAKE A DIFFERENCE

NUEDEXTA provided significant reduction in PBA episodes.

Adverse events were generally mild to moderate.

NUEDEXTA has broad insurance coverage and copay savings for eligible patients.

Onscreen Text and Narrator:

INDICATION and IMPORTANT SAFETY INFORMATION for NUEDEXTA® (dextromethorphan HBr and quinidine sulfate)

INDICATION:

NUEDEXTA is indicated for the treatment of pseudobulbar affect (PBA).



NUEDEXTA®

(dextromethorphan HBr and 20 mg
quinidine sulfate) capsules 10 mg

PBA occurs secondary to a variety of otherwise unrelated neurologic conditions, and is characterized by involuntary, sudden, and frequent episodes of laughing and/or crying. PBA episodes typically occur out of proportion or incongruent to the underlying emotional state. PBA is a specific condition, distinct from other types of emotional lability that may occur in patients with neurologic disease or injury.

IMPORTANT SAFETY INFORMATION:

CONTRAINDICATIONS:

- **Quinidine and Related Drugs:** NUEDEXTA contains quinidine and should not be used concomitantly with other drugs containing quinidine, quinine, or mefloquine.
- **Hypersensitivity:** NUEDEXTA is contraindicated in patients with a history of NUEDEXTA-, quinine-, mefloquine-, or quinidine-induced thrombocytopenia, hepatitis, bone-marrow depression, lupus-like syndrome, or known hypersensitivity to dextromethorphan (e.g., rash, hives).
- **MAOIs:** NUEDEXTA is contraindicated in patients taking monoamine oxidase inhibitors (MAOIs), or in patients who have taken MAOIs within the preceding 14 days, due to the risk of serious and possibly fatal drug interactions, including serotonin syndrome. Allow at least 14 days after stopping NUEDEXTA before starting an MAOI.
- **Cardiovascular:** NUEDEXTA is contraindicated in patients with a prolonged QT interval, congenital long QT syndrome, history suggestive of torsades de pointes, heart failure, patients receiving drugs that both prolong QT interval and are metabolized by CYP2D6 (e.g., thioridazine and pimozide), patients with complete atrioventricular (AV) block without implanted pacemaker, or at high risk of complete AV block.

Thrombocytopenia and Other Hypersensitivity Reactions: Quinidine can cause immune-mediated thrombocytopenia that can be severe or fatal. Non-specific symptoms, such as lightheadedness, chills, fever, nausea, and vomiting, can precede or occur with thrombocytopenia. NUEDEXTA should be discontinued immediately if thrombocytopenia occurs.

Hepatotoxicity: Hepatitis, including granulomatous hepatitis, has been reported in patients receiving quinidine, generally during the first few weeks of therapy. Discontinue immediately if this occurs.

Cardiac Effects: NUEDEXTA causes dose-dependent QTc prolongation. QT prolongation can cause torsades de pointes-type ventricular tachycardia, with the risk increasing as the degree of prolongation increases. When initiating NUEDEXTA in patients at risk for QT prolongation and torsades de pointes, electrocardiographic (ECG) evaluation of QT interval should be conducted at baseline and 3 to 4 hours after the first dose. Some risk factors include use with CYP3A4 inhibitors or drugs that prolong QT interval, electrolyte abnormalities, bradycardia, or left ventricular hypertrophy or dysfunction. If patients taking NUEDEXTA experience symptoms that could indicate the occurrence of cardiac arrhythmias (e.g., syncope or palpitations), NUEDEXTA should be discontinued, and the patient further evaluated.

Concomitant Use of CYP2D6 Substrates: NUEDEXTA inhibits CYP2D6 and may interact with other drugs metabolized by CYP2D6. Adjust dose of CYP2D6 substrates as needed.

Dizziness: NUEDEXTA may cause dizziness. Take precautions to reduce the risk of falls.

NUEDEXTA®

(dextromethorphan HBr and 20 mg
quinidine sulfate) capsules 10 mg

Serotonin Syndrome: Use of NUEDEXTA with selective serotonin reuptake inhibitors (SSRIs) or tricyclic antidepressants increases the risk of "serotonin syndrome."

Anticholinergic Effects of Quinidine: Monitor for worsening in myasthenia gravis.

Adverse Reactions: The most common adverse reactions (incidence of $\geq 3\%$ and two-fold greater than placebo) in patients taking NUEDEXTA are diarrhea, dizziness, cough, vomiting, asthenia, peripheral edema, urinary tract infection, influenza, increased gamma-glutamyltransferase, and flatulence.

These are not all the risks for use of NUEDEXTA.

To report SUSPECTED ADVERSE REACTIONS, contact Otsuka America Pharmaceutical, Inc. at 1-800-438-9927 or FDA at 1-800-FDA-1088 (www.fda.gov/medwatch).

Please see **FULL PRESCRIBING INFORMATION** at nuedextahcp.com

Onscreen Text:

[NUEDEXTA logo]

Onscreen Text:

© 2024 Otsuka America Pharmaceutical, Inc. All rights reserved. NUEDEXTA® is a registered trademark of Avanir Pharmaceuticals, Inc. and used under license by Otsuka America Pharmaceutical, Inc.

July 2024 18US24EBP0033



Otsuka
Otsuka America Pharmaceutical, Inc.